arXiv:1208.1513v1 [math.DS] 7 Aug 2012

TINKER TOY DYNAMICS

LEE DEVILLE AND EUGENE LERMAN

Abstract. We propose a definition of continuous time dynamical systems made up of interacting open subsystems. The interconnections of subsystems are coded by directed graphs. Our main result is that the appropriate maps of graphs called graph fibrations give rise to maps of the appropriate dynamical systems and thereby to conserved quantities.

Contents

1. Introduction 1
2. Definitions and constructions 2
Graphs and manifolds 2
Open systems and their interconnections 5
Interconnections and graphs 6
3. Maps of dynamical systems from graph fibrations 8
References 13

1. Introduction

The idea that symmetries and conservation laws are tightly linked is a well-established meta-principle in mathematics and physics. Noether's principle is probably the best known manifestation of this general meta-principle, but there are others. For example one can regard the orbit types of points in a phase space with symmetry as conserved quantities. This conservation of orbit type plays an important role in equivariant bifurcation theory (see, for example [5]). The interplay between Noether's principle and conservation of orbit type leads to interesting results in symmetric Hamiltonian systems (see, for example [7]). More recently Golubitsky, Stewart and their collaborators found and successfully used a new aspect of the meta-principle that formalizes certain symmetries of ordinary differential equations as groupoid (not group) invariance [3]. They showed that such groupoid symmetries lead to the existence of certain kind of invariant subsystems, the so called polydiagonals.

In this paper we describe another type of symmetry and the attendant conserved quantities that provide a different perspective on the existence of polydiagonals in the so-called "groupoid formalism" of Golubitsky *et al*. Broadly speaking we take the point of view that mathematical objects of interest don't exist by themselves; maps between objects are equally, if not more, important. From this point of view a group symmetry of a dynamical system is a collection of invertible self-maps in the relevant category of dynamical systems. But why look only at the invertible self-maps? In this paper we extract conserved quantities by looking at the arrows in and out of an object in a category. This may seem unusual, but in fact it is not. Given a vector field X on a manifold M it is very desirable but often difficult to find all other dynamical systems (N, Y) and maps $\varphi : N \to M$ so that Y and X are φ-related (that is $D\varphi \circ Y = X \circ \varphi$). In other words it is desirable to finds maps of dynamical systems $\varphi : (N, Y) \to (M, X)$. For instance maps from the circle with the constant vector field $\frac{\partial}{\partial\theta}$ into (M, X) "are" periodic orbits of X. Similarly it is useful to have maps $\psi : (M, X) \to (N, Y)$ out of (M, X). For instance if $\dim N < \dim M$ a map $\psi : (M, X) \to (N, Y)$ may be thought of as a kind of simplification of the dynamical system (M, X) — an "abstraction of (M, X)" in the language of computer science.

Lee DeVille is supported in part by the NSF grant CMG-0934491 .

There is also a different, complementary way to view our results. Consider a continuous time dynamical system made up of interacting subsystems, a network of simpler systems, if you like. Repetition of subsystems is a kind of symmetry, and hence should give rise to some kind of conserved quantities.

We make this idea precise by constructing an appropriate category of networks of continuous time systems. A network in our sense consists of a finite directed graph G with a set of nodes G_0, a *phase space function* $\mathcal{P}$ that assigns to each node of the graph an appropriate phase space (which we take to be a manifold), a family of open systems $\{w_a\}_{a\in G_0}$ (one for each node a of the graph G) consistent in an appropriate way with the structure of the graph, and an interconnection map $\mathscr{I}$ that turns these open systems into a vector field on the product $\prod_{a\in G_0}\mathcal{P}(a)$ of the phase spaces of the nodes. Our main result, Theorem 3.10, shows that graph fibrations (q.v. Definition 3.1) compatible with phase space functions give rise to maps of dynamical systems, hence to conserved quantities.

Our approach to dynamics on networks is complementary to a theory of networks of coupled cells developed by Stewart, Golubitsky and their co-workers [4, 2] and a theory of coupled cell systems of Michael Field [6] as it focuses on maps between networks.

2. Definitions and constructions

Graphs and manifolds. Throughout the paper *graphs* are directed multigraphs, possibly with loops. More precisely, we use the following definition:

2.1. **Definition.** A graph G consists of two finite sets G_1 (of arrows, or edges), G_0 (of nodes, or vertices) and two maps $\mathfrak{s},\mathfrak{t} : G_1 \to G_0$ (source, target):

$$G = \{s, t \colon G_1 \to G_0\}.$$

We write $G = \{G_1 \rightrightarrows G_0\}$.

The set G_1 may be empty, i.e., we may have $G = \{\emptyset \rightrightarrows G_0\}$.

2.2. **Definition.** A map of graphs $\varphi : A \to B$ from a graph A to a graph B is a pair of maps $\varphi_1 : A_1 \to B_1$, $\varphi_0 : A_0 \to B_0$ taking edges of A to edges of B, nodes of A to nodes of B so that for any edge γ of A we have

$$\varphi_0(\mathfrak{s}(\gamma)) = \mathfrak{s}(\varphi_1(\gamma)) \quad \text{and} \quad \varphi_0(\mathfrak{t}(\gamma)) = \mathfrak{t}(\varphi_1(\gamma)).$$

We will usually omit the indices 0 and 1 and write $\varphi(\gamma)$ for $\varphi_1(\gamma)$ and $\varphi(a)$ for $\varphi_0(a)$.

2.3. **Remark.** The collection of (finite directed multi-)graphs and maps of graphs form a category Graph.

In order to construct networks from graphs we need to have a consistent way of assigning manifolds to nodes of our graphs. We formalize this idea by making the collection of graphs with manifolds assigned to vertices into a category Graph/Man.

2.4. **Definition.** The objects of the category Graph/Man are pairs $(G, \mathcal{P})$ where G is a (finite directed multi-)graph and $\mathcal{P} : G_0 \to \mathsf{Man}$ is a function that assigns to each node a of G a manifold $\mathcal{P}(a)$.[1]

A morphism φ from $(G, \mathcal{P})$ to $(G', \mathcal{P}')$ is a map of graphs $\varphi : G \to G'$ with

$$\mathcal{P}' \circ \varphi = \mathcal{P}.$$

2.5. **Remark.** We think of objects $(G, \mathcal{P})$ of Graph/Man as networks of manifolds.

2.6. **Notation.** Given a category $\mathscr{C}$ we denote the opposite category by $\mathscr{C}^{\mathsf{op}}$.

We adhere to the convention that a *contravariant functor* from a category $\mathscr{C}$ to a category $\mathscr{D}$ is a covariant functor

$$F : \mathscr{C}^{\mathsf{op}} \to \mathscr{D}.$$

Then for any morphism $c \xrightarrow{\gamma} c'$ of $\mathscr{C}$ we have $F(c) \xleftarrow{F(\gamma)} F(c')$ in $\mathscr{D}$.

Next we recall the notion of a *product* in a category $\mathscr{C}$. We do this to draw a distinction between categorical and Cartesian products of finite families of manifolds which we will exploit throughout the paper.

---

[1] We think of $\mathcal{P}$ as an assignment of phase spaces to the nodes of the graph G, and for this reason we refer to $\mathcal{P}$ as a *phase space function.*

2.7. **Definition.** A product of a family $\{c_s\}_{s\in S}$ of objects in a category $\mathscr{C}$ indexed by a set S is an object $\prod_{s'\in S} c_{s'}$ of $\mathscr{C}$ together with a family of morphisms $\{\pi_s : \prod_{s'\in S} c_{s'} \to c_s\}_{s\in S}$ with the following universal property: given an object c' of $\mathscr{C}$ and a family of morphisms $\{f_s : c' \to c_s\}_{s\in S}$ there is a unique morphism $f : c' \to \prod_{s\in S} c_s$ with

$$\pi_s \circ f = f_s \quad \text{ for all } s \in S.$$

2.8. **Remark.** If a product exists then it is unique up to a unique isomorphism. This is a standard fact in category theory.

2.9 (Categorical products in the category Man of manifolds)**.**
Observe that in the category Man of (finite dimensional paracompact smooth) manifolds a product $\prod_{s\in S} M_s$ exists for any family $\{M_s\}_{s\in S}$ of manifolds *provided that the indexing set S is finite.* It can be constructed by ordering the elements of S and taking the Cartesian product. That is, by writing $S = \{s_1, \dots, s_n\}$ and setting $\prod_{s\in S} M_s = \prod_{i=1}^n M_{s_i}$, where the right hand side is the Cartesian product. However, for our purposes it is better to have a construction of the product that does not involve a choice of ordering of the indexing set in question. This may be done as follows. Given a family $\{M_s\}_{s\in S}$ of manifolds, denote by $\bigsqcup_{s\in S} M_s$ their disjoint union.[2] Now define

$$\prod_{s\in S} M_s := \{x : S \to \bigsqcup_{s\in S} M_s \mid x(s) \in M_s \text{ for all } s \in S\}.$$

The projection maps $\pi_s : \prod_{s'\in S} M_{s'} \to M_s$ are defined by

$$\pi_s(x) = x(s).$$

We denote $x(s) \in M_s$ by x_s and think of it as s^{th} "coordinate" of an element $x \in \prod_{s\in S} M_s$. Equivalently we may think of elements of the categorical product $\prod_{s\in S} M_s$ as unordered tuples $(x_s)_{s\in S}$ with $x_s \in M_s$.

2.10. **Definition** (total phase space of a network $(G, \mathcal{P})$)**.** For a pair $(G, \mathcal{P})$ consisting of a graph G and an assignment $\mathcal{P} : G_0 \to \mathsf{Man}$, that is, for an object $(G, \mathcal{P})$ of $\mathsf{Graph}/\mathsf{Man}$ we set

$$\mathbb{P}G \simeq \mathbb{P}(G, \mathcal{P}) := \prod_{a\in G_0} \mathcal{P}(a),$$

the categorical product of manifolds attached to the nodes of the graph G by the phase space function $\mathcal{P}$ and call the resulting manifold $\mathbb{P}G$ the total phase space of the network $(G, \mathcal{P})$.

2.11. **Example.** Consider the graph

$G =$ 

Define $\mathcal{P} : G_0 \to \mathsf{Man}$ by $\mathcal{P}(a) = S^2$ (the two sphere) and $\mathcal{P}(b) = S^3$. Then

$$\mathbb{P}(G, \mathcal{P}) = S^2 \times S^3.$$

2.12. **Notation.** If $G = \{\emptyset \rightrightarrows \{a\}\}$ is a graph with one node a and no arrows, we write $G = \{a\}$. Then for any phase space function $\mathcal{P} : G_0 = \{a\} \to \mathsf{Man}$ we abbreviate $\mathbb{P}(\{\emptyset \rightrightarrows \{a\}\}, \mathcal{P} : \{a\} \to \mathsf{Man}) = \mathbb{P}(a, \mathcal{P} : a \to \mathsf{Man})$ as $\mathbb{P}a$.

2.13. **Proposition.** *The assignment*

$$(G, \mathcal{P}) \mapsto \mathbb{P}G := \prod_{a\in G_0} \mathcal{P}(a)$$

of phase spaces to networks extends to a contravariant functor

$$\mathbb{P} : \mathsf{Graph}/\mathsf{Man}^{\mathsf{op}} \to \mathsf{Man}.$$

---

[2] It may be defined by $\bigsqcup_{s\in S} M_s = \bigcup_{s\in S} M_s \times \{s\}$.

Proof. Suppose $\varphi : (G, \mathcal{P}) \to (G', \mathcal{P}')$ is a morphism in $\mathsf{Graph}/\mathsf{Man}$. That is, suppose $\varphi : G \to G'$ is a map of graphs with $\mathcal{P}' \circ \varphi = \mathcal{P}$. We need to define a map of manifolds

$$\mathbb{P}\varphi : \mathbb{P}G' \to \mathbb{P}G.$$

Since by definition $\mathbb{P}G$ is the product $\prod_{a \in G_0} \mathcal{P}(a)$, the universal property of products implies that in order to define $\mathbb{P}\varphi$ it is enough to define a family of maps

$$\left\{(\mathbb{P}\varphi)_a : \mathbb{P}G' \to \mathcal{P}(a)\right\}_{a \in G_0}.$$

For any node a' of G' we have the canonical projection

$$\pi'_{a'} : \mathbb{P}G' \to \mathcal{P}'(a').$$

We therefore define

$$(\mathbb{P}\varphi)_a := \pi'_{\varphi_a} : \mathbb{P}G' \to \mathcal{P}'(\varphi(a)) = \mathcal{P}(a)$$

for all $a \in G_0$. By the universal property of $\mathbb{P}G = \prod_{a \in G_0} \mathcal{P}(a)$ this defines the desired map $\mathbb{P}\varphi : \mathbb{P}G' \to \mathbb{P}G$.

The universal property of products imply that the map $\mathbb{P}$ on morphisms of $\mathsf{Graph}/\mathsf{Man}$ as defined above is actually a functor. That is,

$$\mathbb{P}(\psi \circ \varphi) = \mathbb{P}\varphi \circ \mathbb{P}\psi$$

for any pair (ψ, φ) of composable morphisms in $\mathsf{Graph}/\mathsf{Man}$. □

2.14. **Example.** Suppose G is a graph with two nodes a, b and no edges, G' is a graph with one node $\{c\}$ and no edges, $\mathcal{P}'(c)$ is a manifold M, $\varphi : G \to G'$ is the only possible map of graphs (it sends both nodes to c). $\mathcal{P} : G_0 \to \mathsf{Man}$ is given by $\mathcal{P}(a) = M = \mathcal{P}(b)$ (so that $\mathcal{P}' \circ \varphi = \mathcal{P}$. Then $\mathbb{P}G' \simeq M$,

$$\mathbb{P}G = \{(x_a, x_b) \mid x_a \in \mathcal{P}(a), x_b \in \mathcal{P}(b)\} \simeq M \times M$$

and $\mathbb{P}\varphi : M \to M \times M$ is the unique map with $(\mathbb{P}\varphi(x))_a = x$ and $(\mathbb{P}\varphi(x))_b = x$ for all $x \in \mathbb{P}G'$. Thus $\mathbb{P}\varphi : M \to M \times M$ is the diagonal map $x \mapsto (x, x)$.

2.15. **Example.** Let $(G, \mathcal{P})$, $(G', \mathcal{P}')$ be as in Example 2.14 above and $\psi : (G', \mathcal{P}') \to (G, \mathcal{P})$ be the map that sends the node c to a. Then $\mathbb{P}\psi : \mathbb{P}G \to \mathbb{P}G'$ is the map that sends (x_a, x_b) to x_a.

2.16. **Remark.** If $(G, \mathcal{P})$ is a graph with a phase function, that is, an object of $\mathsf{Graph}/\mathsf{Man}$, and $\varphi : H \to G$ a map of graphs then $\mathcal{P} \circ \varphi : H \to \mathsf{Man}$ is a phase function and $\varphi : (H, \mathcal{P} \circ \varphi) \to (G, \mathcal{P})$ is a morphism in $\mathsf{Graph}/\mathsf{Man}$. We then have a map of manifolds

$$\mathbb{P}\varphi : \mathbb{P}(H, \mathcal{P} \circ \varphi) \to \mathbb{P}(G, \mathcal{P}).$$

Similarly, a commutative diagram

$$\begin{tikzcd} K \arrow[rr, "j"] \arrow[dr, "\psi"'] & & H \arrow[dl, "\varphi"] \\ & G & \end{tikzcd}$$

of maps of graphs and a phase space function $\mathcal{P} : G \to \mathsf{Man}$ give rise to the commutative diagram of maps of manifolds

$$\begin{tikzcd} \mathbb{P}(K, \mathcal{P} \circ \Psi) & & \mathbb{P}(H, \mathcal{P} \circ \varphi) \arrow[ll, "\mathbb{P}j"'] \\ & \mathbb{P}(G, \mathcal{P}) \arrow[ul, "\mathbb{P}\psi"] \arrow[ur, "\mathbb{P}\varphi"'] & \end{tikzcd} .$$

Open systems and their interconnections. Having set up a consistent way of assigning phase spaces to graphs, we now take up continuous time dynamical systems. We start by recalling a definition of an open (control) systems, which is essentially due to Brockett [1].

2.17. **Definition.** A $\mathsf{continuous\ time\ control\ system}$ (or an $\mathsf{open\ system}$) on a region M is a surjective submersion $p : Q \to M$ and a smooth map $F\colon Q \to TM$ so that

$$F(q) \in T_{p(q)}M$$

for all $q \in Q$. That is, the following diagram $\begin{tikzcd} Q \arrow[r, "F"] \arrow[dr, "p"'] & TM \arrow[d, "\pi"] \\ & M \end{tikzcd}$ commutes. Here $\pi : TM \to M$ is the canonical projection.

2.18**.** Given a manifold U of "control variables" we may consider control systems of the form

$$F : M \times U \to TM. \tag{2.19}$$

Here the submersion $p : M \times U \to M$ is given by

$$p(x, u) = x.$$

The collection of all such control systems forms a vector space that we denote by $\mathsf{Control}(M \times U \to M)$:

$$\mathsf{Control}(M \times U \to M) := \{F : M \times U \to TM \mid F(x, u) \in T_xM\}.$$

Now suppose we are given a finite family $\{F_i : M_i \times U_i \to TM_i\}_{i=1}^N$ of control systems and we want to somehow interconnect them to obtain a closed system $\mathscr{I}(F_1, \dots, F_N)$ that is, a vector field, on the product $\prod_i M_i$. What additional data do we need to define the interconnection map

$$\mathscr{I} : \prod_i \mathsf{Control}(M_i \times U_i \to M_i) \to \Gamma\,(T(\prod_i M_i))?$$

The answer is given by the following proposition:

2.20. **Proposition.** *Given a family $\{p_j : M_j \times U_j \to M_j\}_{j=1}^N$ of projections on the first factor and a family of smooth maps $\{s_j : \prod M_i \to M_j \times U_j\}$ so that the diagrams*

$$\begin{tikzcd} M_j \times U_j \arrow[dr, "p_1"] & \\ \prod M_i \arrow[u, "s_j"] \arrow[r, "pr_j"'] & M_j \end{tikzcd}$$

commute for each index j, there is an interconnection map $\mathscr{I}$ making the diagrams

$$\begin{tikzcd} \prod_i \mathsf{Control}(M_i \times U_i \to M_i) \arrow[r, "\mathscr{I}"] \arrow[d] & \Gamma(T(\prod_i M_i)) \arrow[d, "\varpi_j = D(pr_j)\circ -"] \\ \mathsf{Control}(M_j \times U_j \to M_j) \arrow[r, "\mathscr{I}_j"'] & \mathsf{Control}(\prod_i M_i \xrightarrow{pr_j} M_j) \end{tikzcd}$$

commute for each j. The components $\mathscr{I}_j$ of the interconnection map $\mathscr{I}$ are defined by $\mathscr{I}_j(F_j) := F_j \circ s_j$ for all j.

Proof. The space of vector fields $\Gamma(T(\prod_i M_i))$ on the product $\prod_i M_i$ is the product of vector spaces $\mathsf{Control}(\prod_i M_i \to M_j)$:

$$\Gamma(T(\prod_i M_i)) = \prod_j \mathsf{Control}(\prod_i M_i \xrightarrow{pr_j} M_j).$$

In other words a vector field X on the product $\prod_i M_i$ is a tuple $X = (X_1, \dots, X_N)$, where

$$X_j := D(pr_j) \circ X.$$

$(D(pr_j) : T\prod M_i \to TM_j)$ denotes the differential of the canonical projection $pr_j : \prod M_i \to M_j$.) Each component $X_j : \prod_i M_i \to TM_i$ is a control system.

To define a map from a vector space into a product of vector spaces it is enough to define a map into each of the factors. We have canonical projections

$$\pi_j : \prod_i \mathsf{Control}(M_i \times U_i \to M_i) \to \mathsf{Control}(M_j \times U_j \to M_j), \quad j = 1, \dots, N.$$

Consequently to define the interconnection map $\mathscr{I}$ it is enough to define the maps

$$\mathscr{I}_j : \mathsf{Control}(M_j \times U_j \to M_j) \to \mathsf{Control}(\prod_i M_i \xrightarrow{pr_j} M_j).$$

for each index j.

We therefore define the maps $\mathscr{I}_j : \mathsf{Control}(M_j \times U_j \to M_j) \to \mathsf{Control}(\prod_i M_i \xrightarrow{pr_j} M_j)$, $1 \leq j \leq N$, by

$$\mathscr{I}_j(F_j) := F_j \circ s_j.$$

□

2.21. **Remark.** It will be useful for us to remember that the canonical projections

$$\varpi_j : \Gamma(T \prod M_i)) \to \mathsf{Control}(\prod M_i \to M_j)$$

are given by

$$\varpi_j(X) = D(pr_j) \circ X,$$

where $D(pr_j) : T \prod M_i \to TM_j$ are the differential of the canonical projections $pr_j : \prod M_i \to M_j$.

Interconnections and graphs. We next explain how finite directed graphs whose nodes are decorated with phase spaces give rise to interconnection maps. To do this precisely it is useful to have a notion of an *input trees* of a directed graph. Given a graph, an input tree $I(a)$ of a vertex a is roughly, the vertex itself and all of the arrows leading into it. We want to think of this as a graph in its own right, as follows.

2.22. **Definition** (Input tree)**.** Given a vertex a of a graph G we define the *input tree* $I(a)$ to be a graph with the set of vertices $I(a)_0$ given by

$$I(a)_0 := \{a\} \sqcup \mathfrak{t}^{-1}(a);$$

where, as before, the set $\mathfrak{t}^{-1}(a)$ is the set of arrows in G with target a. The set of edges $I(a)_1$ of the input tree is the set of pairs

$$I(a)_1 := \{(a, \gamma) \mid \gamma \in G_1, \ \mathfrak{t}(\gamma) = a\},$$

and the source and target maps $I(a)_1 \rightrightarrows I(a)_0$ are defined by

$$\mathfrak{s}(a, \gamma) = \gamma \quad \text{and} \quad \mathfrak{t}(a, \gamma) = a.$$

In pictures,



2.23. **Example.** Consider the graph $G =$  as in Example 2.11. Then the input tree $I(a)$ is the graph with one node a and no edges: $I(a) = \bullet a$. The input tree $I(b)$ has three nodes and two edges:

$$I(b) =$$



2.24. **Remark.** For each node a of a graph G we have a natural map of graphs

$$\xi = \xi_a : I(a) \to G, \quad \xi_a(a, \gamma) = \gamma,$$

which need not be injective.

2.25**.** Let G be a graph with a phase space function $\mathcal{P} : G \to \mathsf{Man}$ and let a be a node of G. We then have a graph $\{a\}$ with one node and no arrows. Denote the inclusion of $\{a\}$ in G by ι_a and the inclusion into its input tree $I(a)$ by j_a. Then the diagram of graph of maps of graphs

$$\begin{tikzcd}
\{a\} \arrow[rr, "j_a"] \arrow[dr, "\iota_a"'] & & I(a) \arrow[dl, "\xi"] \\
& G &
\end{tikzcd}$$

commutes. By Remark 2.16 we have a commuting diagram of maps of manifolds

$$\begin{tikzcd}
\mathbb{P}\{a\} & & \mathbb{P}I(a) \arrow[ll, "\mathbb{P}j_a"'] \\
& \mathbb{P}G \arrow[ul, "\mathbb{P}\iota_a"] \arrow[ur, "\mathbb{P}\xi"'] &
\end{tikzcd}$$

2.26**.** Let us now examine more closely the map $\mathbb{P}j_a : \mathbb{P}I(a) \to \mathbb{P}a$ in 2.25 above. Since the set of nodes $I(a)_0$ of the input tree $I(a)$ is the disjoint union

$$I(a)_0 = \{a\} \sqcup \mathfrak{t}^{-1}(a),$$

and since $\xi_a(\gamma) = \mathfrak{s}(\gamma)$ for any $\gamma \in \mathfrak{t}^{-1}(a) \subset I(a)_0$, we have

$$\mathbb{P}I(a) = \mathcal{P}(a) \times \prod_{\gamma\in\mathfrak{t}^{-1}(a)} \mathcal{P}(\mathfrak{s}(\gamma)).$$

Since $j_a : \{a\} \to I(a)_0 = \{a\} \sqcup \mathfrak{t}^{-1}(a)$ is the inclusion,

$$\mathbb{P}j_a : \mathbb{P}I(a) \to \mathbb{P}a$$

is the projection

$$\mathcal{P}(a) \times \prod_{\gamma\in\mathfrak{t}^{-1}(a)} \mathcal{P}(\mathfrak{s}(\gamma)) \to \mathbb{P}a.$$

Similarly

$$\mathbb{P}\iota_a : \mathbb{P}G \to \mathbb{P}a$$

is the projection

$$\prod_{b\in G_0} \mathcal{P}(b) \to \mathcal{P}(a).$$

Putting 2.25 and 2.26 together we get

2.27. **Proposition.** *Given a graph G with a phase space function $\mathcal{P} : G_0 \to \mathsf{Man}$, that is, an object $(G, \mathcal{P})$ of $\mathsf{Graph}/\mathsf{Man}$ we have commutative diagrams of maps of manifolds*

$$\begin{tikzcd}
\mathbb{P}I(a) = \mathcal{P}(a) \times \prod_{\gamma\in\mathfrak{t}^{-1}(a)} \mathcal{P}(\mathfrak{s}(\gamma)) \arrow[r, "\mathbb{P}j_a"] & \mathcal{P}(a) \\
\prod_{b\in G_0} \mathcal{P}(b) \arrow[u, "\mathbb{P}\xi"] \arrow[ur, "\mathbb{P}\iota_a"'] &
\end{tikzcd}$$

for each node a of the graph G.

2.28. **Example.** Suppose $G =$  is a graph as in Example 2.11 and suppose $\mathcal{P} : G_0 \to \mathsf{Man}$ is a phase space function. Then

$$\mathbb{P}I(b) \simeq \mathcal{P}(a) \times \mathcal{P}(s) \times \mathcal{P}(b),$$

$\mathbb{P}j_b$ is the projection $\mathcal{P}(a) \times \mathcal{P}(s) \times \mathcal{P}(b) \to \mathcal{P}(b)$ and

$$\mathsf{Control}(\mathbb{P}I(b) \to \mathbb{P}b) = \mathsf{Control}(\mathcal{P}(a) \times \mathcal{P}(s) \times \mathcal{P}(b) \to \mathcal{P}(b)).$$

On the other hand $\mathbb{P}I(a) = \mathcal{P}(a)$, $\mathbb{P}j_a : \mathcal{P}(a) \to \mathcal{P}(a)$ is the identity map and

$$\mathsf{Control}(\mathbb{P}I(a) \to \mathbb{P}a) = \mathscr{X}(\mathcal{P}(a)),$$

the space of vector fields on the manifold $\mathcal{P}(a)$.

2.29. **Notation.** Given $(G, \mathcal{P}) \in \mathsf{Graph}/\mathsf{Man}$ we have a product of vector spaces

$$\mathcal{C}\mathit{trl}(G, \mathcal{P}) := \prod_{a \in G_0} \mathsf{Control}(\mathbb{P}I(a) \to \mathbb{P}a).$$

The elements of $\mathcal{C}\mathit{trl}(G, \mathcal{P})$ are unordered tuples of $(w_a)_{a \in G_0}$ of control systems (q.v. 2.9). We may think of them as sections of the vector bundle $\bigsqcup_{a \in G_0} \mathsf{Control}(\mathbb{P}I(a) \to \mathbb{P}a) \to G_0$ over the vertices of G.

It is easy to see that Propositions 2.20 and 2.27 give us

2.30. **Theorem.** *Given a an object* $(G, \mathcal{P})$ *of the category* $\mathsf{Graph}/\mathsf{Man}$, *there exists a natural interconnection map*

$$\mathscr{I} : \prod_{a \in G_0} \mathsf{Control}(\mathbb{P}I(a) \to \mathbb{P}a) \to \Gamma(T\mathbb{P}G))$$

with

$$\varpi_a \circ \mathscr{I}((w_b)_{b \in G_0}) = w_a \circ \mathbb{P}j_a$$

for all nodes $a \in G_0$. *Here* $\varpi_a : \Gamma T\mathbb{P}G) \to \mathsf{Control}(\mathbb{P}G_0 \xrightarrow{\mathbb{P}\iota_a} \mathbb{P}a)$ *are the projection maps;* $\varpi_a = D(\mathbb{P}\iota_a)$ *(q.v. Remark 2.21).*

2.31. **Example.** Consider the graph $G =$  as in Example 2.11 with a phase space function $\mathcal{P} : G_0 \to \mathsf{Man}$. Then the vector field

$$X = \mathscr{I}(w_a, w_b) : \mathcal{P}(a) \times \mathcal{P}(b) \to T\mathcal{P}(a) \times T\mathcal{P}(b)$$

is of the form

$$X(x, y) = (w_a(x), w_b(x, x, y)) \quad \text{for all } (x, y) \in \mathcal{P}(a) \times \mathcal{P}(b).$$

If $G =$  and $\mathcal{P} : G_0 \to \mathsf{Man}$ is a phase space function, then

$$(\mathscr{I}(w_a, w_b, w_c))\,(x, y, z) = (w_a(x), w_b(x, x, y), w_c(y, z))$$

for all $(w_a, w_b, w_c) \in \mathcal{C}\mathit{trl}(G, \mathcal{P})$ and all $(x, y, z) \in \mathcal{P}(a) \times \mathcal{P}(b) \times \mathcal{P}(c))$.

3. Maps of dynamical systems from graph fibrations

Following Boldi and Vigna [8] (see also [9]) we single out a class of maps of graphs called graph fibrations.

3.1. **Definition.** A map $\varphi : G \to G'$ of directed graphs is a graph fibration if for any vertex a of G and any edge e' of G' ending at $\varphi(a)$ there is a unique edge e of G ending at a with $\varphi(e) = e'$.

3.2. **Example.** The map of graphs



sending the edge γ to γ' and the edge δ to δ' is a graph fibration.

3.3. Given any maps $\varphi : G \to G'$ of graphs and a node a of G there is an induced map of input trees

$$\varphi_a : I(a) \to I(\varphi(a)).$$

On edges of $I(a)$ the map is defined by

$$\varphi(a, \gamma) := (\varphi(a), \varphi(\gamma))$$

(cf. Definition 2.22). Moreover the diagram of graphs

$$\begin{tikzcd}
I(a) \arrow[r, "\varphi_a"] \arrow[d, "\xi_a"'] & I(\varphi(a)) \arrow[d, "\xi_{\varphi(a)}"] \\
G \arrow[r, "\varphi"] & G'
\end{tikzcd}$$

commutes (the map $\xi_a : I(a) \to G$ from an input tree to the original graph is defined in Remark 2.24).

3.4. **Lemma.** *If $\varphi : G \to G'$ is a graph fibration then the induced maps*

$$\varphi_a : I(a) \to I(\varphi(a))$$

of input trees defined above are isomorphisms for all nodes a of G.

Proof. Given an edge $(\varphi(a), \gamma')$ of $I(\varphi(a))$ there is a unique edge γ of G with $\varphi(\gamma) = \gamma'$ and $\mathfrak{t}(\gamma) = a$ and consequently $\varphi_a(a, \gamma) = (\varphi(a), \gamma')$. It follows that φ_a is bijective on vertices and edges. ☐

Recall that a morphism in the category $\mathsf{Graph}/\mathsf{Man}$ from $(G, \mathcal{P})$ to $(G', \mathcal{P}')$ is a map of graphs $\varphi : G \to G'$ with the property that

$$\mathcal{P}' \circ \varphi = \mathcal{P}.$$

3.5. **Definition.** We say that a morphism $\varphi : (G, \mathcal{P}) \to (G', \mathcal{P}')$ in $\mathsf{Graph}/\mathsf{Man}$ is a $\mathsf{graph\ fibration}$ if $\varphi : G \to G'$ is a graph fibration.

The theorem 3.6 below is our reason for singling out graph fibrations.

3.6. **Theorem.** *A graph fibration $\varphi : (G, \mathcal{P}) \to (G', \mathcal{P}')$ in $\mathsf{Graph}/\mathsf{Man}$ naturally induces a linear map*

$$\varphi^* : \mathcal{C}trl(G', \mathcal{P}') \to \mathcal{C}trl(G, \mathcal{P}).$$

Proof. Since

$$\mathcal{C}trl(G, \mathcal{P}) = \prod_{a \in G_0} \mathsf{Control}(\mathbb{P}I(a) \to \mathbb{P}a)$$

is a product of vector spaces, the map φ^* is uniquely determined by maps from $\mathcal{C}trl(G', \mathcal{P}')$ to the factors $\mathsf{Control}(\mathbb{P}I(a) \to \mathbb{P}a)$, $a \in G_0$. On the other hand we have canonical projections

$$\pi_b : \mathcal{C}trl(G', \mathcal{P}') = \prod_{c \in G'_0} \mathsf{Control}(\mathbb{P}I(c) \to \mathbb{P}c) \to \mathsf{Control}(\mathbb{P}I(b) \to \mathbb{P}b)$$

for all $b \in G'_0$. Hence in order to define the map φ^* it is enough to define maps of vector spaces

$$\varphi_a^* : \mathsf{Control}(\mathbb{P}I(\varphi(a)) \to \mathbb{P}\varphi(a)) \to \mathsf{Control}(\mathbb{P}I(a) \to \mathbb{P}a)$$

for all nodes a of the graph G. By 3.3 the diagram

$$\begin{tikzcd}
I(a) \arrow[r, "\varphi_a"] \arrow[d, "\xi_a"'] & I(\varphi(a)) \arrow[d, "\xi_{\varphi(a)}"] \\
G \arrow[r, "\varphi"] \arrow[dr, "\mathcal{P}"'] & G' \arrow[d, "\mathcal{P}'"] \\
& \mathsf{Man}
\end{tikzcd} \tag{3.7}$$

commutes for each $a \in G_0$. Let

$$\varphi|_{\{a\}} : \{a\} \to \{\varphi(a)\}$$

denote the restriction of $\varphi : G \to G'$ to the subgraph $\{a\} \hookrightarrow G$. It is easy to see that the diagrams

$$\begin{tikzcd} I(a) \arrow[r, "\varphi_a"] & I(\varphi(a)) \\ \{a\} \arrow[u, "j_a"] \arrow[r, "\varphi|_{\{a\}}"] & \{\varphi(a)\} \arrow[u, "j_{\varphi(a)}"'] \end{tikzcd} \tag{3.8}$$

commutes as well. By Lemma 3.4 the map φ_a is an isomorphism of graphs. Hence

$$\mathbb{P}\varphi_a : \mathbb{P}I(a) \to \mathbb{P}I(\varphi(a))$$

is an isomorphism of manifolds. Define

$$\varphi_a^* : \mathsf{Control}(\mathbb{P}I(\varphi(a)) \to \mathbb{P}\varphi(a)) \to \mathsf{Control}(\mathbb{P}I(a) \to \mathbb{P}a)$$

by

$$\varphi_a^*(F) = D\mathbb{P}(\varphi|_{\{a\}}) \circ F \circ (\mathbb{P}\varphi_a)^{-1}$$

for all $F \in \mathsf{Control}(\mathbb{P}I(\varphi(a))$. By the universal property of products this gives us the desired map φ^*. Moreover the diagrams

$$\begin{tikzcd} \mathcal{C}\mathit{trl}(G', \mathcal{P}') \arrow[r, "\varphi^*"] \arrow[d, "\pi_{\varphi(a)}"'] & \mathcal{C}\mathit{trl}(G, \mathcal{P}) \arrow[d, "\pi_a"] \\ \mathsf{Control}(\mathbb{P}I(\varphi(a)) \to \mathbb{P}\varphi(a)) \arrow[r, "\varphi_a^*"] & \mathsf{Control}(\mathbb{P}I(a) \to \mathbb{P}a)\} \end{tikzcd}$$

commute for all $a \in G_0$. □

3.9. **Example.** Consider the graph fibration



as in Example 3.2. Let $\mathcal{P}' : G_0' \to \mathsf{Man}$ be a phase space function. Then

$$\mathcal{C}\mathit{trl}(G', \mathcal{P}') = \{(w_a : \mathcal{P}(a) \to T\mathcal{P}(a), w_b : \mathcal{P}'(a) \times \mathcal{P}'(a) \times \mathcal{P}'(b) \to T\mathcal{P}(b), w_c : \mathcal{P}(b) \times \mathcal{P}(c) \to T\mathcal{P}(c))\},$$

$$\mathcal{C}\mathit{trl}(G, \mathcal{P}' \circ \varphi) = \{(w'_{a_1} : \mathcal{P}'(a) \to \mathcal{P}'(a), w_{a_2} : \mathcal{P}'(a) \to \mathcal{P}'(a), w_b : \mathcal{P}'(a) \times \mathcal{P}'(a) \times \mathcal{P}'(b) \to T\mathcal{P}'(b))\}$$

and

$$\varphi^*(w'_a, w'_b, w'_c) = (w'_a, w'_a, w'_b).$$

We are now in the position to state and prove the main result of the paper.

3.10. **Theorem.** *Let $\varphi : (G, \mathcal{P}) \to (G', \mathcal{P}')$ be a graph fibration. Then the pullback map*

$$\varphi^* : \mathcal{C}\mathit{trl}(G', \mathcal{P}') \to \mathcal{C}\mathit{trl}(G, \mathcal{P})$$

constructed in Theorem 3.6 is compatible with the interconnection maps

$$\mathscr{I}' : \mathcal{C}\mathit{trl}(G', \mathcal{P}')) \to \Gamma(T\mathbb{P}G') \quad \textit{and} \quad \mathscr{I} : (\mathcal{C}\mathit{trl}(G, \mathcal{P})) \to \Gamma(T\mathbb{P}G).$$

Namely for any $w' \in \mathcal{C}trl(G', \mathcal{P}')$ *the diagram*

$$\begin{tikzcd}
T\mathbb{P}G' \arrow[r, "D\mathbb{P}\varphi"] & T\mathbb{P}G \\
\mathbb{P}G' \arrow[u, "\mathscr{I}'(w')"] \arrow[r, "\mathbb{P}\varphi"'] & \mathbb{P}G \arrow[u, "\mathscr{I}(\varphi^* w')"']
\end{tikzcd} \tag{3.11}$$

commutes. Consequently

$$\mathbb{P}\varphi : (\mathbb{P}(G', \mathcal{P}'), \mathscr{I}'(w')) \to (\mathbb{P}(G, \mathcal{P}), \mathscr{I}(\varphi^* w'))$$

is a map of dynamical systems.

Proof. Recall that the manifold $\mathbb{P}G$ is the product $\prod_{a \in G_0} \mathbb{P}a$. Hence the tangent bundle bundle $T\mathbb{P}G$ is the product $\prod_{a \in G_0} T\mathbb{P}a$. The canonical projections

$$T\mathbb{P}G \to T\mathbb{P}a$$

are differential of the maps $\mathbb{P}\iota_a : \mathbb{P}G \to \mathbb{P}a$ where, as before, $\iota_a : \{a\} \hookrightarrow G$ is the canonical inclusion of graphs. Hence by the universal property of products, two maps into $T\mathbb{P}G$ are equal if and only if all their components are equal. Therefore, in order to prove that (3.11) commutes it is enough to show that

$$D\mathbb{P}\iota_a \circ \mathscr{I}(\varphi^* w') \circ \mathbb{P}\varphi = D\mathbb{P}\iota_a \circ D\mathbb{P}\varphi \circ \mathscr{I}'(w')$$

for all nodes $a \in G_0$. Note that by definition of the restriction $\varphi|_{\{a\}}$ of $\varphi : G \to G'$ to $\{a\} \hookrightarrow G$, the diagram

$$\begin{tikzcd}
\{a\} \arrow[r, "\varphi|_{\{a\}}"] \arrow[d, "\iota_a"'] & \{\varphi(a)\} \arrow[d, "\iota_{\varphi(a)}"'] \\
G \arrow[r, "\varphi"'] & G'
\end{tikzcd} \tag{3.12}$$

commutes. By the definition of the pullback map φ^* and the interconnection maps $\mathscr{I}$, $\mathscr{I}'$ the diagram

$$\begin{tikzcd}
T\mathbb{P}a & T\mathbb{P}\varphi(a) \arrow[l, "D\mathbb{P}\varphi|_{\{a\}}"] \\
\mathbb{P}I(a) \arrow[u, "(\varphi^* w')_a"'] & \mathbb{P}I(\varphi(a)) \arrow[l, "\mathbb{P}\varphi_a"'] \arrow[u, "w'_{\varphi(a)}"] \\
\mathbb{P}G \arrow[u, "\mathbb{P}\xi_a"'] \arrow[uu, bend left=60, "\mathscr{I}(\varphi^* w')_a"] & \mathbb{P}G' \arrow[l, "\mathbb{P}\varphi"] \arrow[u, "\mathbb{P}\xi_{\varphi(a)}"] \arrow[uu, bend right=60, "\mathscr{I}'(w')_{\varphi(a)}"']
\end{tikzcd} \tag{3.13}$$

commutes as well. We now compute:

$$\begin{aligned}
D\mathbb{P}\iota_a \circ \mathscr{I}(\varphi^* w') \circ \mathbb{P}\varphi &= (\mathscr{I}(\varphi^* w'))_a \circ \mathbb{P}\varphi \\
&= D\mathbb{P}(\varphi|_{\{a\}}) \circ \mathscr{I}'(w')_{\varphi(a)} && \text{by (3.13)} \\
&= D\mathbb{P}(\varphi|_{\{a\}}) \circ D\mathbb{P}\iota_{\varphi(a)} \circ \mathscr{I}'(w') && \text{by definition of } \mathscr{I}'(w')_{\varphi(a)} \\
&= D\mathbb{P}\left(\iota_{\varphi(a)} \circ \varphi|_{\{a\}}\right) \circ \mathscr{I}'(w') && \text{since } \mathbb{P} \text{ is a contravariant functor} \\
&= D\mathbb{P}\left(\varphi \circ \iota_a\right) \circ \mathscr{I}'(w') && \text{by (3.12)} \\
&= D\mathbb{P}(\iota_a) \circ D\mathbb{P}\varphi \circ \mathscr{I}'(w').
\end{aligned}$$

And we are done. ☐

3.14. **Example.** Consider the graph fibration



as in Examples 3.2 and 3.9. Let $\mathcal{P}' : G'_0 \to \mathsf{Man}$ be a phase space function and let $\mathcal{P} = \mathcal{P}' \circ \varphi$. Then

$$\mathbb{P}G' = \mathcal{P}'(a) \times \mathcal{P}'(b) \times \mathcal{P}'(c),$$

$$\mathbb{P}G = \mathcal{P}'(a) \times \mathcal{P}'(a) \times \mathcal{P}'(b),$$

$$\mathbb{P}\varphi(x, y, z) = (x, x, y),$$

and

$$D\mathbb{P}\varphi(p, q, r) = (p, p, q).$$

For any $w' = (w'_a, w'_b, w'_c) \in \mathit{Ctrl}(G', \mathcal{P}')$,

$$\left(\mathscr{I}'(w')\right)(x, y, z) = (w'_a(x), w'_b(x, x, y), w'_c(y, z)),$$

$$\varphi^* w' = (w'_a, w'_a, w'_b),$$

$$\left(\mathscr{I}(\varphi^* w')\right)(x_1, x_2, y) = (w'_a(x_1), w'_a(x_2), w'_b(x_1, x_2, y))$$

and

$$\left(\mathscr{I}(\varphi^* w') \circ \mathbb{P}\varphi\right)(x, y, z) = (w'_a(x), w'_a(x), w'_b(x, x, y))$$

while

$$\left(D\mathbb{P}\varphi \circ \mathscr{I}'(w')\right)(x, , y, z) = D\mathbb{P}\varphi(w'_a(x), w'_b(x, x, y, w'_c(y, z)) = (w'_a(x), w'_a(x), w'_b(x, x, y)).$$

Hence

$$\left(\mathscr{I}(\varphi^* w') \circ \mathbb{P}\varphi\right) = \left(D\mathbb{P}\varphi \circ \mathscr{I}'(w')\right)$$

as expected.

3.15. **Remark.** Theorem 3.10 allows us to define a category DSN of continuous time dynamical systems on networks. The objects of this category are triples

$$(G, \mathcal{P} : G_0 \to \mathsf{Man}, w \in \mathit{Ctrl}(G, \mathcal{P})),$$

where as before G is a finite directed graph, $\mathcal{P}$ is a phase space function and $w = (w_a)_{a \in G_0}$ is a tuple of control systems associated with the input trees of the graph G and the function $\mathcal{P}$.

A morphism from $(G', \mathcal{P}', w')$ to $(G, \mathcal{P}, w)$ is a graph fibration $\varphi : G \to G'$ (yes, the direction of the map is correct) with $\mathcal{P}' \circ \varphi = \mathcal{P}$ and $\varphi^* w' = w$.

Theorem 3.10 allows us to define a functor from DSN to the category DS of continuous time dynamical systems on manifolds. The objects DS are pairs (M, X), where X is a vector field on M. A morphism from (M, X) to (N, Y) in this category is a smooth map $f : M \to N$ with $Df \circ X = Y \circ f$ The functor

$$\mathsf{DSN} \to \mathsf{DS}$$

is defined by

$$\left((G', \mathcal{P}', w') \xrightarrow{\varphi} (G, \mathcal{P}, w)\right) \mapsto (\left(\mathbb{P}G', \mathscr{I}(w')) \xrightarrow{\mathbb{P}\varphi} (\mathbb{P}G, \mathscr{I}(w)\right).$$

Given a dynamical system on a network $(G, \mathcal{P}, w)$ we can forgets all the information except for the graph. This gives rise to a functor

$$\mathsf{DSN} \to \mathsf{FinGraph_{gf}}^{\mathsf{op}}.$$

Here the superscript $^{\mathsf{op}}$ indicates that the functor reverses the direction of arrows and the subscript $_{\mathsf{gf}}$ reminds us that the morphisms in the target category are the (opposite of the) graph fibrations.

References

[1] R.W. Brockett, Control theory and analytical mechanics, in *The 1976 Ames Research Center (NASA) Conference on Geometric Control Theory (Moffett Field, Calif., 1976)*, Math Sci Press, Brookline, Mass., 1977.

[2] Martin Golubitsky, Ian Stewart, and Andrei Török. Patterns of synchrony in coupled cell networks with multiple arrows. *SIAM J. Appl. Dyn. Syst.*, 4(1):78–100 (electronic), 2005.

[3] Martin Golubitsky and Ian Stewart. Nonlinear dynamics of networks: the groupoid formalism. *Bull. Amer. Math. Soc. (N.S.)*, 43(3):305–364, 2006.

[4] Ian Stewart, Martin Golubitsky, and Marcus Pivato. Symmetry groupoids and patterns of synchrony in coupled cell networks. *SIAM J. Appl. Dyn. Syst.*, 2(4):609–646 (electronic), 2003.

[5] M. Field, *Dynamics and symmetry*, ICP Advanced Texts in Mathematics, **3**. Imperial College Press, London, 2007. xiv+478 pp.

[6] M. Field, Combinatorial dynamics, *Dyn. Syst.* **19** (2004), no. 3, 217-243.

[7] J.-P. Ortega and T.S. Ratiu, *Momentum maps and Hamiltonian reduction.* Progress in Mathematics, **222**. Birkhäuser Boston, Inc., Boston, MA, 2004. xxxiv+497 pp.

[8] Paolo Boldi and Sebastiano Vigna. Fibrations of graphs. *Discrete Math.*, 243(1-3):21–66, 2002.

[9] Sebastiano Vigna. `http://vigna.dsi.unimi.it/fibrations/`. Accessed on 9/16/2011.

Department of Mathematics, University of Illinois, Urbana, IL 61801